

# Our Experience with Revascularization and Replantation of Hand

Dr. Yusra Afzal, Dr. Shehab Beg, Dr. Sobia Yasmeen, Dr. Obaid ur Rahman

## ABSTRACT

**Background:** Hand is the main functional unit of body. An amputation is a sum of vascular injury, an open fracture, a soft tissue injury and a nerve injury. Reattachments of amputated parts can result in severe morbidity both during and after surgery. Replantation of a limb continues to be technically demanding procedure. We share our experience of revascularization and replantation of complete and partially amputated hands.

**Material and Methods:** Two year retrospective study, which include patients with complete or incomplete amputation of hand. All patients underwent either revascularization or replantation surgery. Patient's demographics, mode of injury, level of injury, ischemia time and survival were studied. Outcome was assessed on the bases of patients' satisfaction.

**Results:** We studied 16 cases, with amputation of hand. 5 patients had crush injury while 11 faced sharp injury. 4 replantation and 12 revascularization surgeries were performed. The rate of survival was 62.5 %. Successful revascularizations were 66.6 % (8 of 12 patients) and replantations were 75% (3 of 4 patients). All the patients were satisfied with their results

**Conclusion:** The majority of the included patients exhibited good or very good function of the extremity, none of the replanted appendages regained normal levels of functionality. All participants were very satisfied with their outcomes

## Introduction

Hand is the most important part of human body. Hand is a complex combination of different types of bones and joints. Loss of hand or part of it not only effect a man's working ability but it also has an impact on a person's psychology. Traumatic hand amputation is the most catastrophic event in young working individuals<sup>5</sup>. Depression occurs in about  $\geq 30\%$  of amputees<sup>4,8</sup>. Psychological morbidity, decreased self-esteem, and social isolation are also observed in short and long-term follow up after amputation<sup>9,10</sup>.

Replantation surgery is now a standard approach to upper limb amputation around the world<sup>2</sup>. The Quality of life of the patient is preserved by revascularization and replantation of traumatic hand injury<sup>5</sup>. The survival rate of replanted digits is 80 to 90 % in literature<sup>6-7</sup>.

## Materials and Methods

This was a retrospective study conducted at department of plastic surgery at a tertiary care hospital in Karachi .After institutional review board approval, patients with hand amputation who underwent revascularization or replantation from July 2017 to June 2019 were included in the study. All these patients have either complete or incomplete amputation of any part of hand distal to the wrist level. Amputations above wrist joint

*Department of Plastic Surgery,  
Liaquat National Hospital Karachi.  
Corresponding author: Shehab Beg  
Email: shehabbeg@hotmail.com*

and patients who had been operated in any other hospital and arrived for revision surgery were excluded from our study.

We reviewed our data base for revascularization and replantation and assessed it for demographic profile, mode of injury, level of injury, ischemia time, survival and follow up. Outcome was assessed on the bases of patients' satisfaction. Figure 1 shows demography, level of amputation ischemia time and procedure performed

### Procedure

All the patients were assessed in emergency department and were planned. In operating

room, the amputee and stump were assessed and damaged tissue was carefully removed. Bone ends were shortened and rejoined with k wires to hold the bones in place and to allow the rest of the tissues to be restored without tension. Anastomosis of vessels and coaptation of nerves were done. Muscles and tendons were then repaired. In some patients grafts of skin, tendons and blood vessels were needed, too.

All patients had post operative splintage for 4-6 weeks followed by physiotherapy.



**Figure1:** Amputation of thumb & Replantation done

| Sr. No. | Age | Gender | Mechanism of injury | Level of amputation            | Ischemia time at arrival | Procedure |
|---------|-----|--------|---------------------|--------------------------------|--------------------------|-----------|
| 1       | 27  | Male   | Crush               | Thumb Proximal phalynx         | 5 hrs                    | RV        |
| 2       | 31  | Male   | Crush               | CMJ                            | 2 hrs                    | RP        |
| 3       | 25  | Female | Sharp               | Proximal phalylx middle finger | 3.5 hrs                  | RV        |
| 4       | 24  | Male   | Crush               | Mid plalynx index              | 4 hrs                    | RV        |
| 5       | 35  | Male   | Crush               | Wrist joint                    | 5 hrs                    | RV        |
| 6       | 38  | Male   | Sharp               | MPJ                            | 3 hrs                    | RP        |
| 7       | 29  | Male   | Sharp               | Distal phalynx                 | 2 hrs                    | RV        |
| 8       | 33  | Male   | Crush               | Metacarpal                     | 1.5 hrs                  | RP        |
| 9       | 28  | Male   | Crush               | Thumb IPJ                      | 6 hrs                    | RP        |
| 10      | 40  | Male   | Sharp               | Base of metacarpals            | 4 hrs                    | RV        |
| 11      | 48  | Male   | Crush               | Wrist joint                    | 5 hrs                    | RV        |
| 12      | 18  | Male   | Crush               | Index and middle finger        | 6 hrs                    | RV        |
| 13      | 22  | Female | Sharp               | Rt index middle phalnx         | 2.5 hrs                  | RV        |
| 14      | 34  | Male   | Crush               | CMCJ                           | 8 hrs                    | RV        |
| 15      | 36  | Male   | Crush               | Head of metacarpal             | 4 hrs                    | RV        |
| 16      | 38  | Male   | Crush               | CMCJ                           | 5 hrs                    | RV        |

**Table 1:-** Details of injuries to the patients





**Figure 2:** Amputation at wrist & Replantation (Late follow up)

**Results**

We treated 16 patients of hand amputation in two years. Male to female ratio was 7:1. Mean age group was 31.6 years (range 18 to 48 years). Out of 16 patients, our 5 patients (31.3%) had sharp injury and 11 patients (68.7%) had crush injury. 12 patients had revascularization while 4 had replantation surgery

**Outcome**

The rate of survival after revascularization of amputated limb was 66.6 % (8 of 12 patients) and after replantation was 75% (3 of 4 patients). Average of 62.5 %. The outcome of successful hand replantation and revascularization was assessed by subjective patients’ satisfaction. Three month follow up shows that 7 patients had good functional outcome and 3 patients had excellent outcome. However 1 patient was lost to follow up.

|    | <b>Procedure</b>          | <b>Ischemia time</b> | <b>Survival (48 hours)</b> | <b>Outcome 3 months</b> |
|----|---------------------------|----------------------|----------------------------|-------------------------|
| 1  | Thumb revascularization   | 7 hrs                | Viable                     | Good                    |
| 2  | Hand replantation         | 4.5 hrs              | Viable                     | Excellent               |
| 3  | Finger revascularization  | 5 hrs                | Viable                     | Excellent               |
| 4  | Revascularization         | 6.5 hrs              | Viable                     | Good                    |
| 5  | Revascularization         | 7.5 hrs              | Non viable                 | ——                      |
| 6  | Thumb replantation        | 6 hrs                | Viable                     | Good                    |
| 7  | Digit revascularization   | 3.5 hrs              | Viable                     | Excellent               |
| 8  | Hand replantation         | 7 hrs                | Viable                     | Good                    |
| 9  | Replantation              | 8 hrs                | Non viable                 | ——                      |
| 10 | Digit revascularization   | 7 hrs                | Non viable                 | ——                      |
| 11 | Wrist revascularization   | 7 hrs                | Viable                     | Good                    |
| 12 | Finger revascularization  | 8 hrs                | Non viable                 | Good                    |
| 13 | Fingers revascularization | 6 hrs                | Viable                     | Good                    |
| 14 | Hand revascularization    | 10 hrs               | Viable                     | Good                    |
| 15 | Hand revascularization    | 7 hrs                | Non viable                 | Good                    |
| 16 | Hand revascularization    | 8 hrs                | Viable                     | Good                    |

**Table 2:-** Details of Procedures undertaken

## Discussion

In 1962, the first successful limb replantation was performed on a 12 year old boy after traumatic amputation<sup>2</sup>, it was reported by Ronald Malt in 1964<sup>3</sup>. With the passage of time and advances in microvascular surgery, salvage of amputated limb has become a common practice in many centers around the world. Razana et al. described an overall success rate of 65.6%<sup>11</sup> in 1998. Now the success rate has reached upto 80-90%<sup>1</sup>. Our success rate of replantation was 75%, that is at par with other tertiary care unit.

Following the high survival rate of replanted and revascularized post-traumatic hand amputation, recent emphasis has now shifted to functional recovery of the restored part rather than mere survival only. Although the functional outcome of replanted hands will never equal to that of the normal healthy counterpart, the aim of surgery is to produce major functional, cosmetic and psychological improvements in these patients.

In this study, analysis of patient demographic factors (age, sex) and factors related to amputation (level and type of injury) showed no significant influence in the overall survival rate. However, significantly better survival rates were observed in patients who were operated within 6 h compared to those who were operated after 6 h.

## Conclusion

Mode and level of injury and ischemia time are important predictors of success rate of revascularization and replantation of upper limb. Multiple-level, diffuse crush, or avulsion injuries, even if the injuries were incomplete had less return of function.

## References

1. Medling, BD, Bueno, RA, Russell, RC. Replantation outcomes. *Clin Plast Surg* 2007; 34: 177–185, vii–viii.
2. Pao- Yuan Lin, Seng Feng Jeng, Tsan-Shiun Lin. Upper Limb Replantation. *Trauma* oct 2012, 14(4):313-326
3. Malt, RA, McKhann, C. Replantation of severed arms. *JAMA* 1964; 189: 716–722.
4. Engstrom B, Van de Ven, C. *Therapy for Amputees*. 3rd Ed. Churchill Livingstone. 1999
5. T. Kamarul, A Mansor, N. Robson, SHH Albusaidi, AM Suhaeb and Ely Z Samsudin. Replantation and revascularization of amputated upper limb appendages outcome and predicting the factors influencing the success rates of these procedures in a tertiary hospital: An 8-year retrospective, cross-sectional study. *J ortho Surg*. 2018 : 26(1); 1-9
6. Boulas HJ. Amputations of the fingers and hand: indications for replantation. *J Am Acad Orthop Sur*. 1998;6:100–105.
7. Pederson WC. Replantation. *Plast Reconstr Surg*. 2001;107:823–841.
8. Ghous M. Depression: prevalence among Amputees. *Professional Medical Journal*, 2015; 22(2): 263-266
9. Srivastava K, Saldanha D, Chaudhury S, Ryali V, Goyal S, Bhattacharyya D, Basannar D. A Study of Psychological Correlates after Amputation. *Medical Journal Armed Forces India* 2010; 66(4):367-373.
10. Sahu A, Sagar R, Sarkar S, Sagar S. Psychological effects of amputation: A review of studies from India. *Industrial Psychiatry Journal* 2016; 25(1):4–10. doi: 10.4103/0972-6748.196041
11. Razana, A, Hyzan, MY, Pathmanathan, V. Hand replantation and revascularization – six years experience in Hospital Kuala Lumpur 1990–1995. *Med J Malaysia* 1998; 53(Suppl A): 121–130.